


## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup>:</b> <b>C07H 19/06, A61K 31/70</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/14831</b>  <b>(43) International Publication Date:</b> <b>7 July 1994 (07.07.94)</b>
<b>(21) International Application Number:</b> PCT/CA93/00553 <b>(22) International Filing Date:</b> 20 December 1993 (20.12.93)  <b>(30) Priority Data:</b> 07/993,726                   18 December 1992 (18.12.92)   US  <b>(71) Applicant:</b> THE UNIVERSITY OF ALBERTA [CA/CA]; Faculty of Pharmacy and Pharmaceutical Sciences, Edmonton, Alberta T6G 2N8 (CA).  <b>(72) Inventors:</b> KNAUS, Edward, Elmer; 3531-104A Street, Edmonton, Alberta T6J 2K2 (CA). WIEBE, Leonard, Irving; 11022-85 Avenue, Edmonton, Alberta T6G 0W6 (CA). KUMAR, Rakesh; 329-K, Michener Park, Edmonton, Alberta T6H 4M5 (CA).  <b>(74) Agent:</b> JOHNSTON, N., Malcolm, S.; Malcolm Johnston & Associates, Suite 505; 133 Richmond Street West, Toronto, Ontario M5H 2L3 (CA).		<b>(81) Designated States:</b> AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>

**(54) Title:** DIHYDROPYRIMIDINE NUCLEOSIDES WITH ANTIVIRAL PROPERTIES**(57) Abstract**

Pharmaceutical compounds of general formula (I) have been prepared and non-toxic pharmaceutically acceptable salts thereof, wherein R<sub>1</sub> is a halogen substituent; R<sub>2</sub> is a member selected from the group consisting of alkoxy, hydroxy and azido; and X-Y is a member selected from the group consisting of CH(N<sub>3</sub>)-CH<sub>2</sub>,

**(I)**

CH(F)-CH<sub>2</sub> and CH=CH. Halogen denotes an iodo, bromo, chloro and fluoro atom. Alkoxy denotes a straight or branched chain moiety having 1-16 carbon atoms. Compounds of formula (I) can exist as the (5R, 6R), (5S, 6S), (5R, 6S) and (5S, 6R) diastereomers which differ in configuration at positions C-5 and C-6. These compounds exhibit anti-human immunodeficiency virus activity (anti-HIV) and are useful in the treatment of acquired immunodeficiency syndrome (AIDS) and AIDS-related complex.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DIHYDROPYRIMIDINE NUCLEOSIDES WITH ANTIVIRAL PROPERTIES

5

FIELD OF THE INVENTION

The present invention relates to pharmaceutical compounds. More particularly, the invention provides new  
10 unnatural 5,6-dihydropyrimidine nucleoside derivatives, or non-toxic pharmaceutically acceptable salts thereof, having useful physiological antiviral effects, particularly anti-human immunodeficiency virus (anti-HIV) properties which are useful in the treatment of acquired immuno-  
15 deficiency syndrome (AIDS) and AIDS-related complex. The invention relates to such compounds and compositions thereof, and to processes for making and using them.

BACKGROUND OF THE INVENTION

20

Human immunodeficiency virus-1 reverse transcriptase (HIV-1 RT) plays an important role in the life cycle of the virus and has been a major target for the design of drugs to combat AIDS. One class of HIV-1 RT  
25 inhibitors are pyrimidine nucleoside analogs such as 3'-azido-3'-deoxythymidine (AZT), 3'-fluoro-3'-deoxythymidine (FT) and 2',3'-dideohydro-2',3'-dideoxythymidine (d4T). These compounds are converted into their triphosphates by cellular enzymes, the triphosphates  
30 are then recognized by HIV-1 RT as substrates. The corresponding nucleoside monophosphate moiety is incorporated into deoxyribonucleic acid (DNA) chains. Since these analogs lack a 3'-hydroxyl group, this incorporation leads to DNA chain termination. Although AZT appears to be  
35 temporarily effective in decreasing mortality and morbidity in some patients with AIDS, or AIDS-related complex, bone marrow toxicity and anemia are very severe [see the Medical

-2-

Letter, 28 , 107 (1986)]. Frequently administered high doses of AZT must be used to maintain a therapeutic drug level due to its short biological half-life of one hour [see D.D. Richman, M.A. Fischl, M.H. Grieco, M.S. Gottlieb, P.A. Volberding, O.L. Laskin, J.M. Leedom, J. Groopman, D. Mildvan, M.S. Hirsch, G.G. Jackson, D.T. Durack and S. Nusinoff-Lehrman, N. Engl. J. Med., 317, 192 (1987)] which is attributed to its rapid metabolism to the inactive 5'-O-glucuronide (GAZT) and the highly toxic 3'- amino-3'-deoxythymidine (AMT) [see E.M. Cretton, M.-Y. Xie, R.J. Bevan, N.M. Goudgoan, R.F. Schinazi and J.-P. Sommadossi, Mol. Pharmacol., 39 , 258 (1991)]. Since AZT does not penetrate into brain tissue from the cerebral spinal fluid, it does not effectively suppress viral replication in the brain and it is believed that the HIV replicates more rapidly in the central nervous system (CNS), the CNS serving as a reservoir for the virus in the body.

A correlation between lipophilicity, membrane permeability and CNS penetration has long been established [see C. Hansch, A.R. Stewart, S.M. Anderson and D. Bentley, J. Med. Chem., 11 , 1 (1968); D.P. Hall and C.G. Zubrod, Ann. Rev. Pharmacol., 2 , 109 (1962); W.H. Oldendorf, Proc. Soc. Exp. Biol. Med., 147 , 813 (1974)]. The lipophilicity of a compound can be described as the partition coefficient (P) of a drug between 1-octanol (lipid phase) and aqueous buffer at a pH of 7. It has been reported that the partition coefficients for AZT, FT and d4T are 0.964, 0.529 and 0.154, respectively [see E.J. Lien, H. Gao and H. Prabhaker, J. Pharm. Sci., 80 , 517 (1991)]. Although AZT is the most lipophilic, it is neither lipophilic nor hydrophilic since it partitions almost equally (P = 0.964). Several studies to design more lipophilic compounds, and hence their ability to penetrate into the CNS across the blood-brain-barrier (BBB) have not

-3-

resulted so far in compounds with an acceptable therapeutic potency.

5           Although a number of 5,6-dihydrothymidine analogs of the physiological nucleoside thymidine are known (see A.G. Samuel, H.B. Mereyala and K.N. Ganesh, Nucleosides & Nucleotides, 11 , 49 (1992); R. Teoule, B. Fouque and J. Cadet, Nucl. Acid Res., 2 , 487 (1975); G. Bernardinelli, 10 R. Benhamza and J.M. Tronchet, Acta Cryst. C45 , 1917 (1989)) these analogs act as competitive inhibitors of thymidine kinase at low concentrations (see B. Fouque and R. Teoule, Chemotherapy, 20 , 221 (1974)). Since these analogs do not inhibit reverse transcriptase, they are in- 15 effective in the treatment of AIDS or AIDS-related complex.

It has now been discovered that the introduction of a halogen atom in position 5 in conjunction with an alkoxy, hydroxy or azido substituent in position 6 increase 20 lipophilicity thereby resulting in an increased ability to penetrate into the CNS. Such compounds exhibit anti-human immunodeficiency virus (anti-HIV) activity and may also be useful to treat other clinical conditions such as hepatitis B viral infections and other viral infections.

25           In addition such compounds have a longer biological half-life allowing for a longer duration of action and they exhibit an increasing drug stability and a decreasing toxicity. Alternatively, such compounds may serve as pro-drugs, since a reducing agent (such as 30 glutathione in vivo) would regenerate the 5,6-olefinic bond releasing AZT, FT or d4T.


#### DETAILED DESCRIPTION OF THE INVENTION

35           The present invention relates to new 5,6-dihydro-pyrimidine derivatives and non-toxic, pharmaceutically acceptable salts thereof (as well as pharmaceutical

-4-

compositions containing them).

The new compounds according to the present  
5 invention have the general formula:



15

wherein:

$R_1$  is a halogen;  
 $R_2$  is a hydroxy, alkoxy group or azido; and  
 20 X-Y is a member selected from the group CH=CH, CH(N<sub>3</sub>)-CH<sub>2</sub> or CH(F)-CH<sub>2</sub> as well as the non-toxic, pharmaceutically acceptable salts thereof.

The term "halogen" as used herein means fluorine, chlorine, bromine or iodine.

25 The term "alkoxy" as used herein means substituents of straight and branched chain aliphatic alcohols having from 1 to 16 carbon atoms.

Compounds of formula (I) can exist as one of four possible diastereomers wherein  $R_1$  and  $R_2$  have the meanings given above since an asymmetric carbon is respectively  
 30 present at the C-5 and C-6 positions.


The term "diastereomer" means the (5R,6R), (5S,6S), (5R,6S) or (5S,6R) configuration.

The 5-halo-6-alkoxy-5,6-dihydrothymidine

35

-5-

derivatives are prepared by reacting a thymidine analog of the formula:



10 wherein X-Y is a member selected from the group consisting of  $\text{CH}(\text{N}_3)-\text{CH}_2$ ,  $\text{CH}(\text{F})-\text{CH}_2$  and  $\text{CH}=\text{CH}$  with an electrophilic source of halogen of the formula:




wherein  $\text{R}_1$  is an iodo, bromo, chloro or fluoro atom and Z is a member independently selected from the group consisting of iodo, bromo and chloro, in the presence of an alkyl alcohol of the formula:

20



wherein  $\text{R}_2$  is an alkoxy group wherein the alkyl moiety is a straight or branched aliphatic alkyl chain having from 1 to 16 carbon atoms, allowing the reaction to occur in the temperature range of  $-78^\circ\text{C}$  to  $25^\circ\text{C}$ , preferably in the  $0^\circ\text{C}$  to  $25^\circ\text{C}$  range, to convert to 5-halo-6-alkoxy-5,6-dihydrothymidine diastereomers of the formula:

25




wherein  $\text{R}_1$ ,  $\text{R}_2$  and X-Y are as defined above. The reactions are allowed to take place in inert organic

-6-

solvents such as tetrahydrofuran, dioxane or dimethoxyethane when the alkyl alcohol of formula (IV) is a solid.

5 Alternatively, compounds of formula (I) can also be prepared by reacting a thymidine analog of formula (II) wherein X-Y is as defined above, with an electrophilic source of halogen of the formula:

10



(V)

15 wherein R<sub>1</sub> is a member selected from the group consisting of iodo, bromo and chloro, in the presence of an alkyl alcohol of formula (IV) wherein R<sub>2</sub> is as defined as above and glacial acetic acid, allowing the reaction to occur at 25°C to convert to 5-halo-6-alkoxy-5,6-dihydrothymidine derivatives of the formula (I) wherein R<sub>1</sub>, R<sub>2</sub> and X-Y are as defined as above. These reactions are allowed to take place in inert organic solvents such as dimethoxyethane, dioxane or tetrahydrofuran (preferably dimethoxyethane).

25 The 5-halo-6-azido-5,6-dihydrothymidine derivatives are prepared by reacting a thymidine analog of the formula (II) wherein X-Y is as defined as above, with an electrophilic source of halogen of the formula (V) wherein R<sub>1</sub> is as defined above, in an inert organic solvent such as dimethoxyethane, dioxane or tetrahydrofuran, preferably dimethoxyethane, and an alkali metal azide of the formula (VI):

35




(VI)

wherein R<sub>2</sub> is an azido group and M is selected from a group consisting of sodium, lithium and potassium, prefer-



-7-

ably sodium, in a water solvent, allowing the reaction to occur in the -5°C to 25°C range to convert to 5-halo-6-azido-5,6-dihydrothymidine diastereomers of the formula:



wherein  $R_1$  is a member selected from the group consisting of iodo, bromo and chloro,  $R_2$  is an azido substituent and X-Y is as defined above.

The 5-halo-6-hydroxy-5,6-dihydrothymidine derivatives are prepared by reacting a thymidine analog of formula (II) wherein X-Y is as defined above, with an electrophillic source of halogen of the formula (V) wherein  $R_1$  is as defined above, in water as a solvent, allowing the reaction to occur at 0°C to convert to 5-halo-6-hydroxy-5,6-dihydrothymidine diastereomers of the formula (I) wherein  $R_1$  is a member selected from the group consisting of iodo, bromo and chloro,  $R_2$  is a hydroxyl substituent and X-Y is as defined above.

More particularly, the compounds listed in the Examples and in Table I, II, and III have been prepared, and through testing, have been found to have anti-human immunodeficiency virus properties (Table IV).

Suitable pharmaceutically acceptable phosphate forms of these compounds include the 5'-O-monophosphate, 5'-O-diphosphate and 5'-O-triphosphate derivatives.

These compounds can be administered either parentally, as by injection, or orally. As a liquid carrier, a carrier such as water or polyethylene glycol, or other physiologically acceptable solvents or dispersing liquids can be used. For oral administration, either solid or liquid carriers may be used. One commonly used solid

-8-

carrier is gum acacia, but others are also suitable. An operative dosage range is between about 0.01 and 200 mg/kg, preferably between 0.1 and 20 mg/kg.

The following non-limitative examples illustrate some selective methods for producing the compounds according to the present invention, as well as comparative data illustrating the anti-human immunodeficiency virus (anti-HIV) effect of representative compounds according to the present invention.


The starting materials for the preparation of compounds of formula (I), viz the thymidine analogs of formula (II), the electrophilic forms of halogen of formula (III) and formula (V), the alkyl alcohols of formula (IV), and azides of formula (VI) are either known or are conveniently prepared from known starting materials from methods known per se.

The following examples are given for the purpose of illustrating the present invention:

#### Example 1

Preparation of 5-bromo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine:

#### Schematic for Example 1



A freshly prepared solution of methyl hypobromite (bromine in methanol) was added dropwise to a solution of 3'-azido-3'-deoxythymidine (0.2 g, 0.75 mmol) in methanol

-9-

(10 mL) at 25°C with stirring until the light yellow color of the reaction mixture persisted. The reaction was allowed to proceed at 25°C for 20 min prior to neutralization to pH 6 using a solution of methanolic sodium hydroxide. Removal of the solvent in vacuo, dissolution of the residue in methanol (5 mL), adsorption onto silica gel (1 g), removal of the solvent in vacuo, and application of this material to the top of a silica gel column (Merck 7734, 100-200  $\mu$ m particle size) followed by elution with chloroform-methanol (95:5, v/v) afforded a mixture of the diastereomers K-1 and K-2 (0.225 g, 79%) as a viscous oil. Analysis found: C, 34.40; H, 4.27; N, 17.85.  $C_{11}H_{16}BrN_5O_5 \cdot \frac{1}{2} H_2O$  requires: C, 34.12, H, 4.42; N, 18.08. The two diastereomers (5R,6R)-5-bromo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-1) and (5S,6S)-5-bromo-5-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-2) were separated using Whatman PLK5F silica gel plates (1 mm thickness) using chloroform-methanol (95.5, v/v) as development solvent.

Diastereomer K-1:  $[\alpha]_D^{25} = +71.7^\circ$  (c 0.0030, MeOH);  $R_f$  0.61; oil; yield (60 mg, 21%);  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.96 (s, 3H,  $CH_3$ ), 2.32 and 2.68 (two m, 1H each, H-2'), 3.46 (s, 3H,  $OCH_3$ ), 3.80 (m, 1H, H-5'), 3.94 (m, 2H, H-4', H-5''), 4.34 (m, 1H, H-3'), 4.95 (s, 1H, H-6), 5.90 (d,  $J_{1',2'} = 6.0$  Hz, 1H, H-1'), 8.64 (s, 1H, NH, exchanges with deuterium oxide);  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  22.82 ( $CH_3$ ), 37.04 (C-2'), 53.21 (C-5), 57.41 ( $OCH_3$ ), 60.06 (C-3'), 62.12 (C-5'), 84.02 (C-4'), 86.66 (C-1'), 89.16 (C-6), 150.58 (C-2 C=O), 167.10 (C-4 C=O).

Diastereomer K-2:  $[\alpha]_D^{25} = -43.3^\circ$  (c 0.0021, MeOH);  $R_f$  0.63; oil; yield (0.148 g, 52%);  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.98 (s, 3H,  $CH_3$ ), 2.26 and 2.96 (m, 2H, H-2'), 3.60 (s, 3H,  $OCH_3$ ), 2.76 (m, 1H, H-5'), 2.94 (m, 1H, H-5''), 4.02 (m, 1H, H-4'), 4.52 (m, 1H, H-3'), 4.59 (s, 1H, H-6), 5.27 (d,  $J_{1',2'} = 6.0$  Hz, 1H, H-1'), 8.53

-10-

(s, 1H, NH, exchanges with deuterium oxide);  $^{13}\text{C}$  NMR (CDCl<sub>3</sub>)  $\delta$  22.66 (CH<sub>3</sub>), 35.01 (C-2'), 53.34 (C-5), 57.15 (OCH<sub>3</sub>), 61.48 (C-3'), 62.86 (C-5'), 85.05 (C-4'), 92.56 (C-1'), 95.27 (C-6), 150.51 (C-2 C=O), 166.83 (C-4 C=O).

### Example 2

Utilizing the general procedure of Example 1 and starting from the appropriately substituted compounds of the formula (II), of formula (III) and of formula (IV), as represented in the schematic for Example 2, the following compounds of the formula (I) are prepared:

#### Schematic for Example 2







TABLE (1)  
5-halo-6-alkoxy-5,6-dihydrothymidine diastereomers prepared according to Example 2

Chemical Name	 (5R,6R)		 (5S,6S)		 (5S,6R)		X-Y	R <sub>f</sub> <sup>a</sup>	[α] <sub>D</sub> <sup>25</sup> (c, MeOH)	mp, °C
	No.	R <sub>1</sub>	R <sub>2</sub>	No.	R <sub>1</sub>	R <sub>2</sub>				
(5R,6R)-5-bromo-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-3	Br	OEt				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.68	+76.6° (0.0036)	123-125
(5S,6S)-5-bromo-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-4	Br	OEt				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.75	-37.3° (0.0035)	oil
(5R,6R)-5-bromo-6-isopropoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-5	Br	O-1-Pr				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.69	+72.6° (0.0034)	oil
(5R,6R)-5-chloro-6-isopropoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-6	Cl	O-1-Pr				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.69	+70.3° (0.0120)	oil
(5R,6R)-5-bromo-6-(1-octyloxy)-5,6-dihydro-3'-azido-3'-deoxythymidine	K-7	Br	O(CH <sub>2</sub> ) <sub>7</sub> Me				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.81	+41.6° (0.0055)	oil
(5R,6R)-5-chloro-6-(1-octyloxy)-5,6-dihydro-3'-azido-3'-deoxythymidine	K-8	Cl	O(CH <sub>2</sub> ) <sub>7</sub> Me				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.84	+37.7° (0.0048)	oil
(5R,6R)-5-bromo-6-(1-hexadecyloxy)-5,6-dihydro-3'-azido-3'-deoxythymidine	K-9	Br	O(CH <sub>2</sub> ) <sub>15</sub> Me				CH(N <sub>3</sub> )-CH <sub>2</sub>	0.84	+27.6° (0.0085)	oil
(5R,6R)-5-bromo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-10	Br	OMe				CH(F)-CH <sub>2</sub>	0.58	+67.2° (0.0023)	oil
(5S,6S)-5-bromo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-11	Br	OMe				CH(F)-CH <sub>2</sub>	0.70	-72.5° (0.0016)	oil
(5R,6R)-5-bromo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-12	Br	OMe				CH=CH	0.57 <sup>b</sup>	+66.0° (0.0060)	93-95
(5S,6S)-5-bromo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-13	Br	OMe				CH=CH	0.57 <sup>b</sup>	-80.0° (0.0036)	oil
(5R,6R)-5-bromo-6-ethoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-14	Br	OEt				CH=CH	0.61 <sup>c</sup>	ND <sup>d</sup>	oil
(5S,6S)-5-bromo-6-ethoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-15	Br	OEt				CH=CH	0.61 <sup>c</sup>	ND	oil

<sup>a</sup>CHCl<sub>3</sub>/MeOH(9:1,v/v) Whatman 25 mm silica gel thin layer plates

<sup>b</sup>Separated by HPLC using a Whatman Partisil M9 10/25 ODS C-18 reverse phase column using water-methanol as eluant at a flow rate of 2mL/min

<sup>c</sup>Not separated by preparative HPLC

<sup>d</sup>ND=not determined

Example 3

5 Preparation of 5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxythymidine.

Chlorine gas (4.7 g) was bubbled slowly into a suspension of 3'-azido-3'-deoxythymidine (10 g, 37.4 mmol) in 98% ethanol (500 mL) at 0°C with stirring until the  
10 light yellow-green color of the resulting solution persisted. The pH of this solution was adjusted to 6.5 using a solution of sodium hydroxide in ethanol and the mixture was filtered. Removal of the solvent from the filtrate in vacuo and separation of the residue obtained by elution  
15 from a silica gel column using chloroform-methanol (97:3, v/v) as eluent gave (5S,6S)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-17), (5R,6R)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-16), and (5S,6R)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-deoxy-  
20 thymidine (K-18), respectively. Analysis found: C, 41.62; H, 5.20; N, 19.81.  $C_{12}H_{18}ClN_5O_5$  requires: C, 41.44; H, 5.21; N, 20.14.

Diastereomer K-16:  $[\alpha]_D^{25} = +63.0^\circ$  (c 0.019, MeOH);  $R_f$  0.67; mp 118-120°C; yield (8 g, 61.5%);  $^1H$   
25 NMR ( $CDCl_3$ )  $\delta$  1.16 (t,  $J=7$  Hz, 3H,  $OCH_2CH_3$ ), 1.82 (s, 3H, C-5  $CH_3$ ), 2.30 (m, 1H, H-2'), 2.64 (m, 2H, H-2' and 5'-OH which exchanges with deuterium oxide), 3.50-3.98 (m, 5H, H-4', H-5',  $OCH_2CH_3$ ), 4.32 (m, 1H, H-3'), 4.92 (s, 1H, H-6), 5.84 (d,  $J_{1',2'}=6.0$  Hz, 1H, H-1'), 8.30  
30 (s, 1H, NH, exchanges with deuterium oxide);  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  14.93 ( $OCH_2CH_3$ ), 21.76 (C-5  $CH_3$ ), 37.04 (C-2'), 60.10 (C-3'), 60.94 (C-5), 62.20 (C-5'), 65.55 ( $OCH_2CH_3$ ), 84.01 (C-4'), 87.04 (C-1'), 87.92 (C-6), 150.62 (C-2 C=O), 166.62 (C-4 C=O).

35 Diastereomer K-17:  $[\alpha]_D^{25} = -15.3^\circ$  (c 0.028, MeOH);  $R_f$  0.72; oil; yield (0.5 g, 3.7%);  $^1H$  NMR

-13-


(CDCl<sub>3</sub>)  $\delta$  1.10 (t, J=7 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>), 1.68 (s, 3H, C-5 CH<sub>3</sub>), 2.10 (m, 1H, H-2'), 2.78 (m, 1H, H-2''), 3.40-3.92 (m, 5H, H-4', H-5', OCH<sub>2</sub>CH<sub>3</sub>), 4.36 (m, 1H, H-3'), 4.48 (s, 1H, H-6), 5.16 (d, J<sub>1',2'</sub>=6.0 Hz, 1H, H-1'), 9.04 (s, 1H, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  14.72 (OCH<sub>2</sub>CH<sub>3</sub>), 21.58 (C-5 CH<sub>3</sub>), 34.94 (C-2'), 61.09 (C-5), 61.56 (C-3'), 62.96 (C-5'), 65.50 (OCH<sub>2</sub>CH<sub>3</sub>), 85.11 (C-4'), 92.78 (C-1'), 93.86 (C-6), 150.49 (C-2 C=O), 166.25 (C-4 C=O).

Diastereomer K-18: [α]<sub>D</sub><sup>25</sup> = +42.1° (c 0.009, MeOH); R<sub>f</sub> 0.61; oil; yield (3.5 g, 26.7%); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.18 (t, J=7 Hz, 3H, OCH<sub>2</sub>CH<sub>3</sub>), 1.78 (s, 3H, C-5 CH<sub>3</sub>), 2.28 (m, 1H, H-2'), 2.68 (m, 1H, H-2''), 3.20 (br s, 1H, 5'-OH, exchanges with deuterium oxide), 3.60-3.98 (m, 5H, H-4', H-5', OCH<sub>2</sub>CH<sub>3</sub>), 4.36 (m, 1H, H-3'), 4.82 (s, 1H, H-6), 5.64 (d, J<sub>1',2'</sub>=6.0 Hz, 1H, H-1'), 8.80 (br s, 1H, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  14.84 (OCH<sub>2</sub>CH<sub>3</sub>), 25.88 (C-5 CH<sub>3</sub>), 36.98 (C-2'), 60.34 (C-3'), 62.16 (C-5'), 66.67 (OCH<sub>2</sub>CH<sub>3</sub>), 66.98 (C-5), 84.15 (C-4'), 87.88 (C-1'), 89.79 (C-6), 151.10 (C-2 C=O), 167.79 (C-4 C=O).

### Schematic for Example 3

25

Schematic for Example 3



### Example 4

Preparation of 5-chloro-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine.

N-Chlorosuccinimide (0.2 g, 1.5 mmol) was added to a solution of 3'-azido-3'-deoxythymidine (0.2 g, 0.75

-14-

mmol) in methanol (10 mL) and glacial acetic acid (0.6 mL) with stirring and the reaction was allowed to proceed at 25°C for 15 h. At this time additional N-chlorosuccinimide (0.2 g, 1.5 mmol) and glacial acetic acid (0.6 mL) were added and the reaction was allowed to proceed at 25°C for 24 h with stirring prior to neutralization to pH 6.5 using methanolic sodium hydroxide. Removal of the solvent in vacuo gave a residue which was dissolved in chloroform (5 mL), the chloroform solution was washed with cold water (2 x 5 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent was removed in vacuo. The residue obtained was purified by elution from a silica gel column using chloroform-methanol (95:5, v/v) as eluent to yield a mixture of diastereomers (5R,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-19) and (5S,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine (K-20). Analysis found: C, 39.46; H, 4.87. C<sub>11</sub>H<sub>16</sub>ClN<sub>5</sub>O<sub>5</sub> requires: C, 39.58; H, 4.83. The two diastereomers K-19 and K-20 were separated by PTLC using Whatman PLK5F silica gel plates (1 mm thickness) using chloroform-methanol (95:5, v/v) as development solvent.

Diastereomer K-19:  $[\alpha]_D^{25} = +74.7^\circ$  (c 0.0038, MeOH); R<sub>f</sub> 0.57; oil; yield (0.1 g, 40%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.80 (s, 3H, C-5 CH<sub>3</sub>), 2.30 and 2.63 (two m, 1H each, H-2'), 3.46 (s, 3H, OCH<sub>3</sub>), 3.82 (m, 1H, H-5'), 3.96 (m, 2H, H-4', H-5"), 4.32 (m, 1H, H-3'), 4.90 (s, 1H, H-6), 5.92 (d, J<sub>1',2'</sub> = 6.0 Hz, 1H, H-1'), 8.80 (s, 1H, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 21.60 (CH<sub>3</sub>), 36.95 (C-2'), 57.36 (OCH<sub>3</sub>), 60.04 (C-3'), 60.88 (C-5), 62.05 (C-5'), 83.95 (C-4'), 86.39 (C-1'), 88.62 (C-6), 150.66 (C-2 C=O), 166.71 (C-4 C=O).


Diastereomer K-20:  $[\alpha]_D^{25} = +39.3^\circ$  (c 0.0059, MeOH), R<sub>f</sub> 0.54; oil; yield (45 mg, 18%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.83 (s, 3H, C-5 CH<sub>3</sub>), 2.32 and 2.75 (two m,



-15-

1H each, H-2'), 3.56 (s, 3H, OCH<sub>3</sub>), 3.80 (m, 1H, H-5'), 3.98 (m, 2H, H-4', H-5''), 4.40 (m, 1H, H-3'), 4.76 (s, 1H, H-6), 5.78 (d, J<sub>1',2'</sub>=6.0 Hz, 1H, H-1'), 8.28 (br s, 1H, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 26.05 (CH<sub>3</sub>), 37.0 (C-2'), 58.23 (OCH<sub>3</sub>), 60.35 (C-3'), 62.34 (C-5'), 66.88 (C-5), 84.25 (C-4'), 88.18 (C-1'), 91.46 (C-6), 150.57 (C-2 C=O), 167.02 (C-4 C=O).

#### Schematic for Example 4



#### Example 5

20 Preparation of 5-bromo-6-hydroxy-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine.

N-Bromosuccinimide (80 mg, 0.44 mmol) was added in aliquots to a suspension of 2',3'-didehydro-2',3'-dideoxythymidine (0.1 g, 0.44 mmol) in water (5 mL) at 0°C with stirring. The initial yellow color produced upon addition of each aliquot of N-bromosuccinimide disappeared rapidly. After all the N-bromosuccinimide had been added, the reaction mixture was stirred for 20 min at 0°C. Removal of the solvent in vacuo, dissolution of the residue obtained in ethyl acetate (5 mL), adsorption onto silica gel (1 g), removal of the solvent in vacuo and application of this material to the top of a silica gel column followed by elution with chloroform-methanol (96:4, v/v) as eluent afforded (5R,6R)-5-bromo-6-hydroxy-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine (K-32) and (5S,6S)-5-bromo-6-hydroxy-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine (K-33), respectively. Analysis found: C,

-16-


37.89; H, 4.15; N, 8.63.  $C_{10}H_{13}BrN_2O_5$  requires: C, 37.40; H, 4.07; N, 8.72.

5 Diastereomer K-32:  $[\alpha]_D^{25} = +31.9^\circ$  (c. 0.0026, MeOH);  $R_f$  0.42; mp 94-95°C; yield (60 mg, 43%);  $^1H$  NMR ( $CD_3OD$ )  $\delta$  1.88 (s, 3H,  $CH_3$ ), 3.74 (m, 2H, H-5'), 4.80 (m, 1H, H-4'), 5.15 (s, 1H, H-6), 5.90 (m, 1H, H-3'), 6.30 (m, 1H, H-2'), 6.82 (m, 1H, H-1');  $^{13}C$  NMR ( $CD_3OD$ )  $\delta$  23.38 ( $CH_3$ ), 55.29 (C-5), 62.56 (C-5'), 81.76 (C-6), 87.38 (C-4'), 91.77 (C-1'), 127.14 (C-2'), 135.35 (C-3').

Diastereomer K-33:  $[\alpha]_D^{25} = -32.7^\circ$  (c. 0.0011, MeOH),  $R_f$  0.35; oil; yield (47 mg, 33.1%);  $^1H$  NMR ( $CD_3OD$ )  $\delta$  1.82 (s, 3H,  $CH_3$ ), 3.74 (m, 2H, H-5'), 4.75 (m, 1H, H-4'), 5.28 (s, 1H, H-6), 5.95 (m, 1H, H-3'), 6.24 (m, 1H, H-2'), 6.78 (m, 1H, H-1');  $^{13}C$  NMR ( $CD_3OD$ )  $\delta$  23.30 ( $CH_3$ ), 54.68 (C-5), 65.07 (C-5'), 80.12 (C-6), 87.71 (C-4'), 90.79 (C-1'), 127.80 (C-2'), 133.96 (C-3'), 152.87 (C-2 C=O), 169.92 (C-4 C=O).

#### Schematic for Example 5

#### Schematic for Example 5




#### 30 Example 6

Illustrates the preparation of 5-halo-6-alkoxy-5,6-dihydrothymidines following the alternate method of preparation seen in example 5 and described in the

35 schematic for Example 6.

Schematic for Example 6

5

Schematic for Example 6

15 Starting from the appropriately substituted compounds of formula (II), of formula (IV) and of formula (V), the following compounds of the formula (I) are prepared:

20

25

30

35

TABLE (II)  
5-halo-6-alkoxy-5,6-dihydrothymidine diastereomers prepared according to Example 6

Chemical Name	No	R <sub>1</sub>	R <sub>2</sub>	X-Y	R <sub>f</sub> <sup>a</sup>	[α] <sub>D</sub> <sup>25</sup> (c, MeOH)	mp. °C
(5R, 6R) -5-iodo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-21	1	OMe	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.57	+87.3° (0.0055)	oil
(5S, 6S) -5-iodo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine	K-22	1	OMe	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.63	-46.2° (0.0037)	oil
(5R, 6R) -5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-23	1	OMe	CH(F)-CH <sub>2</sub>	0.61	+56.6° (0.0017)	oil
(5S, 6S) -5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-24	1	OMe	CH(F)-CH <sub>2</sub>	ND <sup>b</sup>	ND	oil
(5S, 6R) -5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-25	1	OMe	CH(F)-CH <sub>2</sub>	0.55	+32.2° (0.0016)	oil
(5R, 6R) -5-iodo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-26	1	OMe	CH(F)-CH <sub>2</sub>	0.58	+74.1° (0.0014)	oil
(5S, 6S) -5-iodo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-27	1	OMe	CH(F)-CH <sub>2</sub>	0.66	-83.0° (0.0035)	oil
(5R, 6R) -5-chloro-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-28	1	OMe	CH=CH	0.52 <sup>c</sup>	+75.5° (0.0041)	138-139
(5S, 6S) -5-chloro-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-29	1	OMe	CH=CH	0.52 <sup>c</sup>	ND	oil
(5R, 6R) -5-iodo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-30	1	OMe	CH=CH	0.54 <sup>c</sup>	ND	oil
(5S, 6S) -5-iodo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-31	1	OMe	CH=CH	0.54 <sup>c</sup>	ND	oil

<sup>a</sup>CHCl<sub>3</sub>/MeOH(9:1,v/v) Whatman 25 OM silica gel thin layer plates

<sup>b</sup>ND=not determined

<sup>c</sup>Separated by HPLC using a Whatman Partisil M9 10/25 ODS, C-18 reverse phase column using water-methanol as eluant at a flow rate of 2mL/min

SUBSTITUTE SHEET

Example 7

5                   Preparation of 5-bromo-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine.

                  N-Bromosuccinimide (36 mg, 2 mmol) was added in aliquots to a precooled (-5°C) suspension prepared by mixing a solution of 3'-azido-3'-deoxythymidine (52 mg, 2  
10 mmol) in dimethoxyethane (10 mL) and a solution of sodium azide (52 mg, 8 mmol) in water (0.125 mL) with stirring. The initial yellow color produced upon addition of each aliquot of N-bromosuccinimide quickly disappeared. When all the N-bromosuccinimide had reacted, the reaction  
15 mixture was stirred for 30 min at 0°C, poured onto ice-water (25 mL) and extracted with ethyl acetate (3 X 50 mL). Washing the ethyl acetate extract with cold water (10 mL), drying the ethyl acetate solution (Na<sub>2</sub>SO<sub>4</sub>) and removal of the solvent in vacuo gave a residue which was separated  
20 by silica gel column chromatography using chloroform as eluent to give a mixture of diastereomers (5R,6R)-5-bromo-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine (K-34), (5S,6S)-5-bromo-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine (K-35), and (5R,6S)-5-bromo-6-azido-5,6-  
25 dihydro-3'-azido-3'-deoxythymidine (K-36), respectively. Analysis found: C, 30.69; H, 3.95; N, 28.77. C<sub>10</sub>H<sub>13</sub>BrN<sub>8</sub>O<sub>4</sub> requires: C, 30.86; H, 3.36; N, 28.79.

                  Diastereomers K-34 and K-35: R<sub>f</sub> 0.63; yield (30  
30 mg, 38.6%); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.98 and 2.0 (two s, 3H total, CH<sub>3</sub>), 2.30-2.74 (m, 2H total, H-2'), 2.94 (br s, 1H, 5'-OH, exchanges with deuterium oxide), 3.82-4.02 (m, 3H total, H-4' and H-5'), 4.30 and 4.36 (two m, 1H total, H-3'), 5.42 and 5.64 (two s, 1H total, H-6), 5.76 and 6.20  
35 (two d, J<sub>1',2'</sub>=6.0 Hz, 1H total, H-1'), 8.60 and 8.68 (two s, 1H total, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 22.76 and 23.08 (CH<sub>3</sub>), 35.99 and


-20-

36.71 (C-2'), 52.31 and 52.79 (C-5), 60.04 and 60.52 (C-6),  
61.72 and 62.35 (C-5'), 73.88 and 76.64 (C-3'), 83.78 and  
5 84.22 (C-4'), 87.81 (C-1'), 149.88 and 150.02 (C-2 C=O),  
166.11 (C-4 C=O).

Diastereomer K-36:  $[\alpha]_D^{25} = -47.5^\circ$  (c.  
0.0016, MeOH);  $R_f$  0.61; yield (20 mg, 25.7%);  $^1H$  NMR  
( $CDCl_3$ )  $\delta$  1.98 (s, 3H,  $CH_3$ ), 2.24 and 2.34 (two m, 1H  
10 each, H-2'), 3.82-4.05 (m, 3H, H-4', H-5'), 4.37 (m, 1H,  
H-3'), 5.74 (s, 1H, H-6), 6.04 (d,  $J_{1',2'} = 6.0$  Hz,  
1H, H-1'), 8.25 (s, 1H, NH, exchanges with deuterium  
oxide);  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  27.63 ( $CH_3$ ), 36.02  
(C-2'), 60.98 (C-6), 61.75 (C-5), 62.65 (C-5'), 74.75  
15 (C-3'), 83.56 (C-4'), 85.05 (C-1'), 149.66 (C-2 C=O),  
166.26 (C-4 C=O).

#### Schematic for Example 7

##### Schematic for Example 7



#### Example 8

Illustrates the preparation of 5-halo-6-azido-  
5,6- dihydrothymidines using a procedure similar to the one  
30 outlined in Example 7. Starting from the appropriately  
substituted compounds of formula (II), formula (V) and  
formula (VI), the following compounds are prepared:

35

TABLE (III)  
5-halo-6-azido-5,6-dihydro-3'-deoxythymidine diastereomers prepared according to Example 8

Chemical Name	No	P <sub>1</sub>	R <sub>2</sub>	X-Y	R <sub>f</sub> <sup>a</sup>	[α] <sub>D</sub> <sup>25</sup> (c, MeOH)	mp, °C
(5R, 6R) - 5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine	K-37	Cl	N <sub>3</sub>	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.63 <sup>b</sup>	ND <sup>c</sup>	oil
(5S, 6S) - 5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine	K-38	Cl	N <sub>3</sub>	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.63 <sup>b</sup>	ND	oil
(5S, 6R) - 5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine	K-39	Cl	N <sub>3</sub>	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.63 <sup>b</sup>	ND	oil
(5R, 6S) - 5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine	K-40	Cl	N <sub>3</sub>	CH(N <sub>3</sub> )-CH <sub>2</sub>	0.63 <sup>b</sup>	ND	oil
(5R, 6R) - 5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-41	Br	N <sub>3</sub>	CH(F)-CH <sub>2</sub>	0.57 <sup>b</sup>	ND	oil
(5S, 6S) - 5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-42	Br	N <sub>3</sub>	CH(F)-CH <sub>2</sub>	0.57 <sup>b</sup>	ND	oil
(5R, 6S) - 5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine	K-43	Br	N <sub>3</sub>	CH(F)-CH <sub>2</sub>	0.57 <sup>b</sup>	ND	oil
(5R, 6R) - 5-bromo-6-azido-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-44	Br	N <sub>3</sub>	CH=CH	0.57 <sup>b</sup>	ND	oil
(5S, 6S) - 5-bromo-6-azido-5,6-dihydro-2',3'-didehydro-2',3'-deoxythymidine	K-45	Br	N <sub>3</sub>	CH=CH	0.57 <sup>b</sup>	ND	oil

<sup>a</sup>CHCl<sub>3</sub>/MeOH(9:1,v/v) Whatman 25 OM silica gel thin layer plates

<sup>b</sup>Not separated by preparative HPLC

<sup>c</sup>ND=not determined

-22-

The compounds listed in the Examples, Tables I, II, and III have been found to have anti-human immunodeficiency virus properties.

1. Anti-human immunodeficiency activity.

The test is designed to measure the efficacy against HIV for drugs acting at any stage of the virus reproductive cycle and involves the killing of T4 lymphocytes by HIV.

In order to test the activity of the compounds according to the invention, all tests were compared with at least one positive (e.g. AZT-treated) control done at the same time under identical conditions.

The test drug is dissolved in dimethylsulfoxide, then diluted 1:100 in cell culture medium before preparing serial half-log<sub>10</sub> dilutions. T4 lymphocytes (CEM cell line) are added and after a brief interval HIV-1 is added, resulting in a 1:200 final dilution of the test drug. Uninfected cells with the test drug serve as a toxicity control, and infected and uninfected cells without the test drug serve as basic controls. Cultures are incubated at 37°C in a 5% CO<sub>2</sub> atmosphere for 6 days. The tetrazolium salt, XTT, is added to all wells, and cultures are incubated to allow formazan color development by viable cells. Individual wells are analyzed spectrophotometrically to quantitate formazan production, and in addition are viewed microscopically for detection of viable cells and confirmation of protective activity. Test drug-treated virus-infected cells are compared with test drug-treated non-infected cells and with other appropriate controls (untreated infected and untreated noninfected cells, test drug-containing wells without cells, etc.) on the same plate [see O.W. Weislow, R. Kiser, D. Fine, J. Bader, R.H. Shoemaker, M.R. Boyd, J. Natl. Cancer Inst., 81, 577 (1989)]. The test results are shown in the following Table IV, the compounds listed being comparable to 3'-azido-3'-



TABLE (IV)

Anti-HIV activity of 5-halo-6-alkoxy (or azido)-5,6-dihydrothymidine diastereomers tested

Substance	IC <sub>50</sub> (M) <sup>a</sup>	EC <sub>50</sub> (M) <sup>b</sup>	TI(IC <sub>50</sub> /EC <sub>50</sub> ) <sup>c</sup>
K-1	1.72 x 10 <sup>-5</sup>	3.27 x 10 <sup>-9</sup>	5260
K-2	4.25 x 10 <sup>-5</sup>	2.80 x 10 <sup>-7</sup>	152
K-3	1.85 x 10 <sup>-5</sup>	6.75 x 10 <sup>-9</sup>	2740
K-4	2.22 x 10 <sup>-5</sup>	2.37 x 10 <sup>-8</sup>	936
K-10	1.72 x 10 <sup>-6</sup>	5.25 x 10 <sup>-9</sup>	328
K-11	9.72 x 10 <sup>-6</sup>	3.25 x 10 <sup>-9</sup>	2991
K-12/K-13 <sup>d</sup>	>1.28 x 10 <sup>-4</sup>	5.46 x 10 <sup>-5</sup>	2
K-14/K-15 <sup>d</sup>	>1.40 x 10 <sup>-5</sup>	ND <sup>e</sup>	ND
K-19/K-20 <sup>d</sup>	>8.98 x 10 <sup>-4</sup>	5.79 x 10 <sup>-6</sup>	155
K-21	1.87 x 10 <sup>-5</sup>	3.17 x 10 <sup>-9</sup>	5899
K-22	6.42 x 10 <sup>-6</sup>	5.15 x 10 <sup>-9</sup>	1247
K-23	>8.0 x 10 <sup>-6</sup>	5.55 x 10 <sup>-9</sup>	144
K-25	>8.0 x 10 <sup>-6</sup>	3.79 x 10 <sup>-9</sup>	21
K-26	5.73 x 10 <sup>-5</sup>	ND	ND
K-27	1.22 x 10 <sup>-5</sup>	3.75 x 10 <sup>-9</sup>	3253
K-28/K-29 <sup>d</sup>	>1.03 x 10 <sup>-3</sup>	3.75 x 10 <sup>-4</sup>	2
K-30/K-31 <sup>d</sup>	6.60 x 10 <sup>-5</sup>	3.75 x 10 <sup>-7</sup>	178
K-32	>2.0 x 10 <sup>-4</sup>	ND	ND
K-33	2.0 x 10 <sup>-4</sup>	ND	ND
K-34/K-35 <sup>d</sup>	1.76 x 10 <sup>-4</sup>	ND	ND
K-37/K-38/K-39/K-40 <sup>d</sup>	3.5 x 10 <sup>-4</sup>	1.49 x 10 <sup>-6</sup>	235
K-41/K-42/K-43 <sup>d</sup>	1.0 x 10 <sup>-4</sup>	1.45 x 10 <sup>-8</sup>	6896
K-44/K-45 <sup>d</sup>	4.47 x 10 <sup>-5</sup>	9.18 x 10 <sup>-7</sup>	49
AZT	5 X 10 <sup>-4</sup>	3 x 10 <sup>-9</sup>	

<sup>a</sup>The IC<sub>50</sub> value is the test drug concentration which results in a 50% survival of uninfected control cells (eg. cytotoxic activity of the test drug)

<sup>b</sup>The EC<sub>50</sub> value is the test drug concentration which produces a 50% survival of HIV infected cells relative to uninfected controls (eg. in vitro anti-HIV activity)

<sup>c</sup>Therapeutic index


<sup>d</sup>Tested as a mixture of diastereomers

<sup>e</sup>ND = not determined

**SUBSTITUTE SHEET**

We claim:

1. A dihydrothymidine derivative of the formula (I):



or a non-toxic pharmaceutically acceptable salt thereof, wherein  $R_1$  is a halogen substituent selected from the group consisting of iodo, bromo, chloro and fluoro;  $R_2$  is a member selected from the group consisting of alkoxy wherein the alkyl moiety is a straight or branched chain having from 1 to 16 carbon atoms, hydroxy and azido; and  $X-Y$  is a member selected from the group consisting of  $CH(N_3)-CH_2$ ,  $CH(F)-CH_2$  and  $CH=CH$ .

- 20 2. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is a methoxy.
3. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is an ethoxy.
- 25 4. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is an isopropoxy.
5. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is a 1-octyloxy.
- 30 6. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is a 1-hexadecyloxy.
7. A dihydrothymidine derivative according to Claim 1, wherein  $R_2$  is a hydroxy or an azido.

-25-

8. (5R,6R)-5-bromo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
- 5
9. (5S,6S)-5-bromo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
10. (5R,6R)-5-bromo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 2.
- 10
11. (5S,6S)-5-bromo-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 2.
- 15
12. (5R,6R)-5-bromo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine according to Claim 2.
13. (5S,6S)-5-bromo-6-methoxy-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine according to Claim 2.
- 20
14. (5R,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
15. (5S,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
- 25
16. (5R,6R)-5-iodo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
- 30
17. (5S,6S)-5-iodo-6-methoxy-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 2.
18. (5R,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 2.
- 35
19. (5S,6S)-5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 2.

-26-

20. (5S,6R)-5-chloro-6-methoxy-5,6-dihydro-3'-fluoro-3'-  
5 deoxythymidine according to Claim 2.
21. (5R,6R)-5-iodo-6-methoxy-5,6-dihydro-3'-fluoro-3'-  
deoxythymidine according to Claim 2.
- 10 22. (5S,6S)-5-iodo-6-methoxy-5,6-dihydro-3'-fluoro-3'-  
deoxythymidine according to Claim 2.
23. (5R,6R)-5-chloro-6-methoxy-5,6-dihydro-2',3'-  
didehydro-2',3'-dideoxythymidine according to Claim 2.
- 15 24. (5S,6S)-5-chloro-6-methoxy-5,6-dihydro-2',3'-  
didehydro-2',3'-dideoxythymidine according to Claim 2.
25. (5R,6R)-5-iodo-6-methoxy-5,6-dihydro-2',3'-  
20 didehydro-2',3'-dideoxythymidine according to Claim 2.
26. (5S,6S)-5-iodo-6-methoxy-5,6-dihydro-2',3'-  
didehydro-2',3'-dideoxythymidine according to claim 2.
- 25 27. (5R,6R)-5-bromo-6-ethoxy-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 3.
28. (5S,6S)-5-bromo-6-ethoxy-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 3.
- 30 29. (5R,6R)-5-bromo-6-ethoxy-5,6-dihydro-2',3'-  
didehydro-2',3'-dideoxythymidine according to Claim 3.
30. (5S,6S)-5-bromo-6-ethoxy-5,6-dihydro-2',3'-  
35 didehydro-2',3'-dideoxythymidine according to Claim 3.
31. (5R,6R)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-

-27-

deoxythymidine according to Claim 3.

5 32. (5S,6S)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 3.

33. (5S,6R)-5-chloro-6-ethoxy-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 3.

10

34. (5R,6R)-5-bromo-6-isopropoxy-5,6-dihydro-3'-azido-  
3'-deoxythymidine according to Claim 4.

35. (5R,6R)-5-chloro-6-isopropoxy-5,6-dihydro-3'-azido-  
15 3'-deoxythymidine according to Claim 4.

36. (5R,6R)-5-bromo-6-(1-octyloxy)-5,6-dihydro-3'-azido-  
3'-deoxythymidine according to Claim 5.

20 37. (5R,6R)-5-chloro-6-(1-octyloxy)-5,6-dihydro-3'-  
azido-3'-deoxythymidine according to Claim 5.

38. (5R,6R)-5-bromo-6-(1-hexadecyloxy)-5,6-dihydro-3'-  
azido-3'-deoxythymidine according to Claim 6.

25

39. (5R,6R)-5-bromo-6-hydroxy-5,6-dihydro-2',3'-  
didehydro-2',3'-dideoxythymidine according to Claim 7.

40. (5S,6S)-5-bromo-6-hydroxy-5,6-dihydro-2',3'-  
30 didehydro-2',3'-dideoxythymidine according to Claim 7.

41. (5R,6R)-5-bromo-6-azido-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 7.

35 42. (5S,6S)-5-bromo-6-azido-5,6-dihydro-3'-azido-3'-  
deoxythymidine according to Claim 7.

-28-


43. (5R,6S)-5-bromo-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 7.
- 5 44. (5R,6R)-5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 7.
- 10 45. (5S,6S)-5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 7.
46. (5S,6R)-5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 7.
- 15 47. (5R,6S)-5-chloro-6-azido-5,6-dihydro-3'-azido-3'-deoxythymidine according to Claim 7.
48. (5R,6R)-5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 7.
- 20 49. (5S,6S)-5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 7.
50. (5R,6S)-5-bromo-6-azido-5,6-dihydro-3'-fluoro-3'-deoxythymidine according to Claim 7.
- 25 51. (5R,6R)-5-bromo-6-azido-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine according to Claim 7.
- 30 52. (5S,6S)-5-bromo-6-azido-5,6-dihydro-2',3'-didehydro-2',3'-dideoxythymidine according to Claim 7.
53. A method of preparing 5-halo-6-alkoxy-5,6-dihydro-

35

-29-

thymidine derivatives of formula (I) as in Claim 2 or 3 or 4 or 5 or 6:

5




10

wherein  $R_1$  is a iodo, bromo, chloro or fluoro atom;  $R_2$  represents a  $C_1$ - $C_{16}$  alkoxy group with a straight or branched alkyl chain and X-Z is  $CH=CH$ ,  $CH(N_3)-CH_2$  or  $CH(F)-CH_2$  which comprises:

15

reacting a thymidine compound of formula (II):

20



25 (III)

with an electrophilic source of halogen of formula



wherein  $R_1$  is as defined above and Z is independently a iodo, bromo or chloro atom.

In the presence of an alkyl alcohol of the formula

30 (IV):



wherein  $R_2$  is as defined above.


54. A method of preparing dihydrothymidine derivatives

35

-30-

according to Claim 53, wherein the electrophilic source of halogen is:

5




(V)

10 wherein R<sub>1</sub> is a iodo, bromo or chloro atom.

55. A method of preparing 5-halo-6-azido-5,6-dihydro-thymidine derivatives of formula (I) as in claim 7:

15




(I)

20

wherein R<sub>1</sub> is a iodo, bromo, chloro or fluoro atom; R<sub>2</sub> is an azido group; and X-Z is CH=CH, CH(N<sub>3</sub>)-CH<sub>2</sub> or CH(F)-C which comprises:

25

reacting a thymidine of formula (II)



(II)

30

with an electrophilic source of halogen of formula (III):



wherein Z is independently a iodo, bromo or chloro atom and  
35 R<sub>1</sub> is as defined above.

In the presence of an alkali metal azide of the




-31-

formula (VI):

$$R_2-M$$


5 wherein  $R_2$  is as defined above and M is a sodium, selected from the group of sodium, lithium and potassium.

56. A method of preparing 5-halo-6-hydroxy-5,6-dihydro-thymidine derivatives of formula (I) as in claim 7:



wherein  $R_1$  is a iodo, bromo, chloro or fluoro atom;  $R_2$  is a hydroxy radical; and X-Z is  $CH=CH$ ,  $CH(N_3)-CH_2$  or  $CH(F)-C$  which comprises:

reacting a thymidine of formula (II):



with an electrophilic source of halogen of formula (III):

$$R_1-Z$$

wherein Z is independently a iodo, bromo or chloro atom and  $R_1$  is as defined above.

In the presence of a solvent of formula (IV):

$$R_2-H$$

wherein  $R_2$  is an hydroxy group.

35

## A. CLASSIFICATION OF SUBJECT MATTER

IPC5: C07H 19/06, A61K 31/70

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC5: C07H, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, CA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Acta Cryst., Volume C49, 1993, H. Blanchard et al, "Structure of (+)-(5R,6R)-5-Chloro-6-methoxy-5, 6-dihydro-1-(2',3'-didehydro-2', 3'-dideoxy-beta-D-glycero-2-enopentofuranosyl)thymi ne" page 246 - page 250.	1-2,23-24, 53-54
Y	SCIENCE, Volume 249, Sept 1990, H. Mitsuya et al, "Molecular Targets for AIDS Therapy" page 1533 - page 1544	1-56



Further documents are listed in the continuation of Box C.



See patent family annex.

## \* Special categories of cited documents

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

17 March 1994

07 04 94

Name and mailing address of the International Searching Authority/Authorized officer



European Patent Office, P.B. 5818 Patentlaan 2  
NL-2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
Fax: (+31-70) 340-3016

EVA JOHANSSON

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	STN International, File CA, Chemical Abstracts, volume 81, no. 25, 23 December 1974 (Columbus Ohio, US), B. Fouque et al: "Inhibition of Ehrlich ascites cell thymidine kinase by a new class of nucleoside derivatives", abstract no. 165382m, & Chemotherapy (Basel), 20(4), 221-6 1974  --	1-56
A	STN International, File CA, Chemical Abstracts, volume 115, no. 11, 16 September 1991 (Columbus, Ohio, US), Lien Eric J. et al: "Physical factors contributing to the partition coefficient and retention time of 2'3'-dideoxynucleoside analogs", J. Pharm. Sci., 80(6), 517-21  --	1-56
A	STN International, File CA, Chemical Abstracts, volume 114, no. 21, 27 May 1991 (Columbus, Ohio, US), Cretton Erika M. et al: "Catabolism of 3'-azido-3'-deoxythymidine in hepatocytes and liver microsomes, with evidence of formation of 3'-amino-3'-deoxythymidine, a highly toxic catabolite for human bone marrow cells", Mol. Pharmacol., 39(2), 258-66  --	1-56
A	STN International, File CA, Chemical Abstracts, volume 110, no. 1, 2 January 1989 (Columbus, Ohio, US), Chu Chung K. et al: "Comparative activity of 2',3'-saturated and unsaturated pyrimidine and purine nucleosides against human immunodeficiency virus type 1 in peripheral blood mononuclear cells", Biochem. Pharmacol., 37(19), 3543-8  --	1-56
A	STN International, File CA, Chemical Abstracts, volume 110, no. 13, 27 March 1989 (Columbus, Ohio, US), Chu Chung K. et al: "Structure-activity relationships of pyrimidine nucleosides as antiviral agents for human immunodeficiency virus type 1 in peripheral blood mononuclear cells", J. Med. Chem., 32(3), 612-17  -----	1-56